arXiv:1608.01363v1 [math.RA] 3 Aug 2016

ON $\mathfrak{F}$-HYPERCENTRAL MODULES AND CHARACTER CLUSTERS

DONALD W. BARNES

Abstract. Let $\mathfrak{F}$ be a saturated formation of soluble Lie algebras over a field F of characteristic $p > 0$ and let $\mathbb{F}_p$ denote the field of p elements. Let $(L, [p])$ be a restricted Lie algebra over F with $z^{[p]} = 0$ for all z in the centre of L. Let $S \in \mathfrak{F}$, $S \neq 0$ be a subnormal subalgebra of L. Let V, W be L-modules. Suppose that the character cluster of W is contained in the set of $\mathbb{F}_p$-linear combinations of the characters in the character cluster of V. Suppose that V, regarded as S-module, is $\mathfrak{F}$-hypercentral. Then W, regarded as S-module, is also $\mathfrak{F}$-hypercentral.

Let F be a field of characteristic $p > 0$ and let $\mathfrak{F}$ be a saturated formation of soluble Lie algebras over F. Let $(L, [p])$ be a restricted Lie algebra over F with $z^{[p]} = 0$ for all z in the centre of L. Suppose that $S \in \mathfrak{F}$, $S \neq 0$ is a subnormal subalgebra of L. If the L-module V is a $[p]$-module, then by [2, Theorem 6.4], V is $S\mathfrak{F}$-hypercentral, that is, $\mathfrak{F}$-hypercentral as S-module. As an L-module is a $[p]$-module if and only if it has character 0, this establishes a link between characters and $\mathfrak{F}$-centrality.

An $S\mathfrak{F}$-hypercentral L-module need not have character 0. Not every L-module has a character. To cope with this situation, the concept of the character cluster $\mathrm{Cl}(V)$ of V was defined in [3, Section 2] to be the set of characters of the composition factors of the module $\bar{V}$ obtained by extending the field to its algebraic closure $\bar{F}$. Since for any saturated formation $\mathfrak{F}$, the trivial 1-dimensional module is $S\mathfrak{F}$-central and has character 0, the following theorem is a generalisation of the theorem cited above. We denote the field of p elements by $\mathbb{F}_p$ and the set of $\mathbb{F}_p$-linear combinations of the elements of $\mathrm{Cl}(V)$ by $\mathbb{F}_p \mathrm{Cl}(V)$.

Theorem 1. *Let $\mathfrak{F}$ be a saturated formation of soluble Lie algebras over a field F of characteristic $p > 0$. Let $(L, [p])$ be a restricted Lie algebra over F with $z^{[p]} = 0$ for all z in the centre of L. Suppose that $S \in \mathfrak{F}$, $S \neq 0$ is a subnormal subalgebra of L. Let V, W be L-modules with $\mathrm{Cl}(W) \subseteq \mathbb{F}_p \mathrm{Cl}(V)$ and suppose that V is $S\mathfrak{F}$-hypercentral. Then W is $S\mathfrak{F}$-hypercentral.*

Proof. Suppose that $\mathrm{Cl}(V)$ has k elements. Put $X = \oplus_{r=1}^{k(p-1)} V^{\otimes r}$. From [5, Theorem 5.2.7(3)], it follows that $\mathrm{Cl}(V^{\otimes r}) = \{c_1 + \cdots + c_r \mid c_i \in \mathrm{Cl}(V)\}$ and so, that $\mathrm{Cl}(X) = \mathbb{F}_p \mathrm{Cl}(V)$. By [1, Theorem 2.1], X is $S\mathfrak{F}$-hypercentral. By replacing V with X, we may suppose that $\mathrm{Cl}(W) \subseteq \mathrm{Cl}(V)$. We need only consider the case where W is irreducible.

Suppose that the character $c \in \mathrm{Cl}(W)$. Then also $c \in \mathrm{Cl}(V)$. Denote the action of $x \in L$ on $\mathrm{Hom}(V, W)$ by $\rho(x)$. By [5, Theorem 5.2.7(1)], $0 = c - c \in \mathrm{Cl}(\mathrm{Hom}(V, W))$. Thus the set $\{(\rho(x)^p - \rho(x^{[p]}))f = 0 \mid x \in L\}$ of linear equations

2010 *Mathematics Subject Classification.* Primary 17B30, 17B50.
Key words and phrases. modular Lie algebras, saturated formations.

over F has a non-zero solution for f in $\bar{F}\otimes\mathrm{Hom}(V,W)$, so it has a non-zero solution in $\mathrm{Hom}(V,W)$. Thus $\mathrm{Hom}(V,W)$ has a non-zero submodule H with character 0. The evaluation map $\mathrm{Eval} : V\otimes H\to W$ given by $\mathrm{Eval}(v,f)=f(v)$ has non-zero image. As Eval is an L-module homomorphism and W is irreducible, we have $\mathrm{Eval}(V\otimes H)=W$. By [2, Theorem 6.4], H is an $S\mathfrak{F}$-hypercentral L-module. As V is $S\mathfrak{F}$-hypercentral, by [1, Theorem 2.1], $V\otimes H$ is $S\mathfrak{F}$-hypercentral. It follows that W is $S\mathfrak{F}$-hypercentral. □

Suppose that we have $S\in\mathfrak{F}$, $S\neq 0$ and S-modules V,W with V $\mathfrak{F}$-hypercentral. We can take any p-envelope $(L,[p])$ of S and, if necessary, adjust the p-operation to have $z^{[p]}=0$ for all z in the centre of L. We then have that S is an ideal of L. Using the construction in the proof of [5, Proposition 2.5.6], we can extend the actions of S on V,W making them into L-modules. (This extension of the action is set out in detail in [4, Section 4].) Suppose that for some choice of $(L,[p])$ and some choices of the extensions of the actions, we have that $\mathrm{Cl}(W)\subseteq\mathbb{F}_p\,\mathrm{Cl}(V)$. Then it follows by Theorem 1, that W also is $\mathfrak{F}$-hypercentral.

References

1. D. W. Barnes, *On $\mathfrak{F}$-hypercentral modules for Lie algebras*, Archiv der Math. **30** (1978), 1–7.
2. D. W. Barnes, *Ado-Iwasawa extras*, J. Aust. Math. Soc. **78** (2005), 407–421.
3. D. W. Barnes, *Character clusters for Lie algebra modules over a field of non-zero characteristic,* Bull. Aust. Math. Soc. **89** (2014), 234–242. Also arXiv:1301.4710.
4. D. W. Barnes, *Induced modules for modular Lie algebras,v.4,* arXiv:1312.0163, (2016).
5. H. Strade and R. Farnsteiner, *Modular Lie algebras and their representations*, Marcel Dekker, Inc., New York-Basel, 1988.

1 Little Wonga Rd., Cremorne NSW 2090, Australia,
E-mail address: D.Barnes@maths.usyd.edu.au